

## Area Under the Bell Curve

In addition to exotic but familiar functions like  $\ln x$ , we can also use definite integrals and Riemann sums to get truly *new* functions.

**Example:** The solution to  $y' = e^{-x^2}$ ;  $y(0) = 0$  is:

$$F(x) = \int_0^x e^{-t^2} dt$$

The graph of  $e^{-x^2}$  is known as the bell curve, and  $F(x)$  describes the area under the curve. This function is extremely useful for computing probabilities.




Figure 1: Graph of  $e^{-x^2}$ .

The exciting thing about  $F(x)$  is that although we have a geometric definition and can compute it using Riemann sums, we can't describe it in terms of any function we've seen previously, including logarithmic and trigonometric functions. It's a completely new function. The problem of describing  $F$  is analogous to the problem of calculating the value of  $\pi$  — the area of a circle with radius 1. The number  $\pi$  is transcendental; it is not the root (zero) of an algebraic equation with rational coefficients.

Using definite integrals we can define a huge class of new functions, many of which are important tools in science and engineering.

MIT OpenCourseWare  
<http://ocw.mit.edu>

18.01SC Single Variable Calculus  
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.